

1. Rationality and consequences

Roberto Lucchetti

Politecnico di Milano

Summary of the slides

- 1 Description of the game
- 2 Players are egoistic
- 3 Rationality assumptions
- 4 Preferences
- 5 Utility functions
- 6 Probability laws
- 7 Deepness of analysis
- 8 Decision theory
- 9 Dominated strategies
- 10 First consequences

Optimization

- 1 One decision maker
- 2 At least two decision makers

Possible variants with one decision maker:

- 1 scalar optimization
- 2 vector optimization
- 3 deterministic optimization
- 4 stochastic optimization
- 5 ...

Many possible variants with many decision makers

- 1 Game theory
- 2 Social choice
- 3 Mechanism design
- 4 Machine learning
- 5 ...

Crucial difference: the best to do is easily definable when there is one decision maker, much more difficult when many decision makers

Description of the game

A process that can be described by:

- 1 A set of players (with more than one element)
- 2 An initial situation
- 3 The way the players must act and all their available moves
- 4 All possible final situations
- 5 The preferences of all agents on the set of the final situations

Examples:

- 1 The chess game
- 2 Two people bargaining how to divide a pie
- 3 A burglar and a guard
- 4 Parties in a Parliament
- 5 ...

Games are **efficient models** for an enormous amount of everyday life situations

Assumptions of the theory.

Players are

- 1 Egoistic
- 2 Rational

Egoistic means that the player cares only about **her own** preferences on the outcomes of the game

This is **not** an ethical issue, but a **mathematical assumption**, aimed at correctly define what means to make a rational choice

Rationality is a much more involved issue

Definition

Let X be a set. A binary relation \succeq on a set X is called:

- reflexive, if for each $x \in X$, xRx
- transitive, if for each $x, y, z \in X$, xRy and $yRz \Rightarrow xRz$
- total, if for each $x, y \in X$, $x \neq y \Rightarrow xRy$ or yRx
- antisymmetric, if for each $x, y \in X$, xRy and $yRx \Rightarrow x = y$

A reflexive, transitive and total binary relation on X is called a **total preorder** (also called, a **ranking**) on X . A reflexive, transitive, total and antisymmetric binary relation on X is called a **total order** on X .

The **first rationality assumption** reads:

The agents are able to provide a preference relation over the outcomes of the game.

Definition

Let \succeq be a preference relation over X . A **utility function** representing \succeq is a function $u : X \rightarrow \mathbb{R}$ such that

$$u(x) \geq u(y) \iff x \succeq y.$$

- 1 A utility function need not to exist, however it exists in general setting, in particular if X is a finite set
- 2 When a utility function exists, then infinite utility functions do exist, since any strictly increasing transformation of a utility function is still a utility function

The **second rationality assumption** reads:

The agents are able to provide a utility function representing their preferences relations, whenever necessary

Allais experiment 1

First shop

Alternative A

gain	probability
2500	33%
2400	66%
0	1%

Alternative B:

gain	probability
2500	0%
2400	100%
0	0%

In a sample of 72 people exposed to this experiment, 82% of them decided to play the Lottery B.

This is rational if $\frac{34}{100} u(2400) > \frac{33}{100} u(2500)$.

Second shop

Alternative C

gain	probability
2500	33%
0	67%

Alternative D:

gain	probability
2400	34%
0	66%

83% of the people interviewed selected lottery C.

This is rational if $\frac{34}{100} u(2400) < \frac{33}{100} u(2500)$.

Thus it is $A \Leftrightarrow C$: Allais experiment shows that usually agents are not rational players!

The **third rationality assumption** reads:

The players use consistently the probability laws, in particular they are consistent w.r.t the calculation of expected utilities, they are able to update probabilities according to Bayes rule...

The beauty contest

Write an integer between 1 and 100.

The mean M is calculated.

Those writing the number at the minimum distance from qM win the game ($0 < q < 1$).

The player imagined by the theory will answer 1 for every q , with little chance to win.

The fourth rationality assumption reads:

The players are able to understand consequences of all actions, consequences of this information on any other player, consequences of the consequences...

Finally, the **fifth rationality assumption** reads:

The player are able to use decision theory, whenever it is possible

This means that, given a set of alternatives X , and a utility function u on X , the player seeks an $\bar{x} \in X$ such that

$$u(\bar{x}) \geq u(x), \forall x \in X.$$

Summarizing the rationality assumptions

- 1 The players are able to rank the outcomes of the game
- 2 The players are able to provide a utility function for their ranking
- 3 The players apply the expected value principle to built their utility function in presence of random events
- 4 The players are able to analyze all consequences of their actions, and the consequences of the consequences and so on
- 5 The players use the apparatus of decision theory anytime it is possible

A first concrete consequence of the axioms

A basic consequence of the “decision theory” assumption is:

A player does not take an action a if she has available an action b providing her a strictly better result, **no matter what the other players do**

Principle of elimination of strictly dominated actions.

Player one action set is $\{18, \dots, 30\}$, player two action set is $\{\text{accept}, \text{refuse}\}$. If player two preference is passing the exam with any grade, rather than repeating it, the action *refuse* is strictly dominated. Please note: asking for raising the grade by one or two points is not an available action

An example

Player 1 chooses a row, player 2 a column. The obtained item is a pair, the first (second) digit is the utility of Player 1 (2):

$$\begin{pmatrix} (8, 8) & (2, 7) \\ (7, 2) & (0, 0) \end{pmatrix}$$

Utilities of player 1:

$$\begin{pmatrix} 8 & 2 \\ 7 & 0 \end{pmatrix}$$

The second row is **strictly dominated** by the first, thus player 1 will select the first row

Even if this principle is usually not very informative, it has surprising consequences

Better Argentina or Italy?

Comparisons of games

The first:

$$\begin{pmatrix} (10, 10) & (3, 15) \\ (15, 3) & (5, 5) \end{pmatrix}$$

The second one:

$$\begin{pmatrix} (8, 8) & (2, 7) \\ (7, 2) & (0, 0) \end{pmatrix}$$

Observe: in any outcome the players are **better off** in the first game rather than in the second:

However it is **more convenient** for them to play the second!

Less is better than more

The first game:

$$\begin{pmatrix} (10, 10) & (3, 5) \\ (5, 3) & (1, 1) \end{pmatrix}$$

The second game, containing all possible outcomes the first, and some further outcomes:

$$\begin{pmatrix} (1, 1) & (11, 0) & (4, 0) \\ (0, 11) & (10, 10) & (3, 5) \\ (0, 4) & (5, 3) & (1, 1) \end{pmatrix}$$

Having less available actions can make the players **better off!**

$$\begin{pmatrix} (0,0) & (1,1) \\ (1,1) & (0,0) \end{pmatrix}$$

Rational outcomes of this game?

We formally do not know but it is obvious that the rational outcomes will be (1,1)

(First row, second column) and (second row, first column) **cannot be distinguished** and this creates a coordination problem between the players

Elimination of dominated strategies

A votation. Three players, alternatives A, B, C . Players preferences:

$$A \succ_{\neq 1}^1 B \succ_{\neq 1} C$$

$$B \succ_{\neq 2} C \succ_{\neq 2} A$$

$$C \succ_{\neq 3} A \succ_{\neq 3} B$$

In case of three different votes, the alternative selected by player one is winning

What can we expect as rational outcome of the game?

Try with elimination of dominated actions. . .

¹ $A \succ_{\neq} B$ means $A \succ B$ and not $B \succ A$

The voting game

- Alternative A is a weakly dominant strategy for Player 1
- Players 2 and 3 have as weakly dominated strategy to play their worst choice

Thus the game reduces to

A	A
C	A

The result is the worst one for the first player!