

Cooperative games (2)

Roberto Lucchetti

Luiss

Contents of the week

Cooperative
games (2)

Roberto
Lucchetti

The nucleolus

The Shapley
value

Power indices

Semivalues

- The nucleolus
- The Shapley value
- Other values
- Examples

Excess

Cooperative
games (2)

Roberto
Lucchetti

The nucleolus

The Shapley
value

Power indices

Semivalues

A TU game v is given

Definition

The **excess** of a coalition A over the imputation x is

$$e(A, x) = v(A) - \sum_{i \in A} x_i$$

$e(A, x)$ is a measure of the **dissatisfaction** of the coalition A with respect to the assignment of the imputation x

Remark

An imputation x of the game v belongs to $C(v)$ if and only if $e(A, x) \leq 0$ for all A

Definition

The **lexicographic** vector attached to the imputation x is the $(2^n - 1)$ -th dimensional vector $\theta(x)$ such that

- ① $\theta_i(x) = e(A, x)$, for some $A \subseteq N$
- ② $\theta_1(x) \geq \theta_2(x) \geq \dots \geq \theta_{2^n-1}(x)$

Definition

The **nucleolus** solution is the solution $\nu : \mathcal{G}(N) \rightarrow \mathbb{R}^n$ such that $\nu(v)$ is the set of the imputations x such that $\theta(x) \leq_L \theta(y)$, for all y imputations of the game v

Remark

$x \leq_L y$ if either $x = y$ or there exists $j \geq 1$ such that $x_i = y_i$ for all $i < j$, and $x_j < y_j$. \leq_L defines a total order in any Euclidean space

An example

Cooperative
games (2)

Roberto
Lucchetti

The nucleolus

The Shapley
value

Power indices

Semivalues

Example

Three players, $v(A) = 1$ if $|A| \geq 2$, 0 otherwise.

Suppose $x = (a, b, 1 - a - b)$, with $a, b \geq 0$ and $a + b \leq 1$. The coalitions S complaining ($e(S, x) > 0$) are those with two members.

$$e(\{1, 2\}) = 1 - (a + b), e(\{1, 3\}) = b, e(\{2, 3\}) = a$$

We must minimize

$$\max\{1 - a - b, b, a\}$$

$$\nu = (1/3, 1/3, 1/3)$$

Remember $C(\nu) = \emptyset$

Nucleolus: one point solution

Cooperative
games (2)

Roberto
Lucchetti

The nucleolus

The Shapley
value

Power indices

Semivalues

Theorem

For every TU game v with nonempty imputation set, the nucleolus $\nu(v)$ is a singleton

Thus the nucleolus is a **one point solution**

Nucleolus in the core

Cooperative
games (2)

Roberto
Lucchetti

The nucleolus

The Shapley
value

Power indices

Semivalues

Proposition

Suppose v is such that $C(v) \neq \emptyset$. Then $\nu(v) \in C(v)$

Proof Take $x \in C(v)$. Then $\theta_1(x) \leq 0$. Thus $\theta_1(\nu)(v) \leq 0$. This implies $\theta_i(\nu)(v) \leq 0$ for all i , i.e. no coalition complains about $\nu(v)$. Then $\nu(v) \in C(v)$ ■

Another example

Cooperative
games (2)

Roberto
Lucchetti

The nucleolus

The Shapley
value

Power indices

Semivalues

$$v(\{1\}) = a, v(\{2\}) = v(\{3\}) = v(\{2, 3\}) = 0, v(\{1, 2\}) = b, v(\{1, 3\}) = c, v(N) = c$$

$$C(v) = \{(x, 0, c - x) : b \leq x \leq c\}$$

Must find x : $\nu(v) = (x, 0, c - x)$. The relevant excesses are

$$e(\{1, 2\}) = b - x, \quad e(\{2, 3\}) = x - c$$

Thus

$$\nu(v) = \left\{ \left(\frac{b+c}{2}, 0, \frac{c-b}{2} \right) \right\}$$

The Shapley value

Cooperative
games (2)

Roberto
Lucchetti

The nucleolus

The Shapley
value

Power indices

Semivalues

Definition

Let v be a TU game. Define the solution assigning the following quantity $\sigma_i(v)$ to player i :

$$\sigma_i(v) = \sum_{S \in 2^N \setminus \{i\}} \frac{s!(n-s-1)!}{n!} [v(S \cup \{i\}) - v(S)]$$

Comments

Cooperative
games (2)

Roberto
Lucchetti

The nucleolus

The Shapley
value

Power indices

Semivalues

The term

$$m_i(v, S) := v(S \cup \{i\}) - v(S)$$

is called the **marginal contribution of player i to coalition $S \cup \{i\}$**

The Shapley value is a weighted sum of all marginal contributions of the players.

Interpretation of the weights

Suppose the players plan to meet in a certain room at a fixed hour, and suppose the **expected arrival time** is the same for all players. If player i enters into the coalition S if and only at her arrival she find in the room **all members of S and only them**, the probability to join coalition S is

$$\frac{s!(n-s-1)!}{n!}$$

An example

Cooperative
games (2)

Roberto
Lucchetti

The nucleolus

The Shapley
value

Power indices

Semivalues

Example

The game:

$$v(\{1\}) = 0, v(\{2\}) = v(\{3\}) = 1, v(\{1, 2\}) = 4, v(\{1, 3\}) = 4, v(\{2, 3\}) = 2, v(N) = 8$$

	1	2	3
123	0	4	4
132	0	4	4
213	3	1	4
231	6	1	1
312	3	4	1
321	6	1	1
	$\frac{18}{6}$	$\frac{15}{6}$	$\frac{15}{6}$

$$\sigma_1(v) = \frac{1!1!}{3!} [v(\{1, 2\}) - v(\{2\})] + \frac{1}{6} [v(\{1, 3\}) - v(\{3\})] + \frac{1}{3} [v(\{N\}) - v(\{2, 3\})] = 3$$

$$\sigma_2(v) = \frac{2}{6} + \frac{5}{6} + \frac{4}{3} = \frac{15}{6}$$

$$\sigma_3(v) = \frac{2}{6} + \frac{5}{6} + \frac{4}{3} = \frac{15}{6}$$

A simple airport game

Cooperative
games (2)

Roberto
Lucchetti

The nucleolus

The Shapley
value

Power indices

Semivalues

Example

The game:

$$v(\{1\}) = 0, v(\{2\}) = v(\{3\}) = 1, v(\{1, 2\}) = 4, v(\{1, 3\}) = 4, v(\{2, 3\}) = 2, v(N) = 8$$

	1	2	3
123	c_1	$c_2 - c_1$	$c_3 - c_2$
132	c_1	0	$c_3 - c_1$
213	0	c_2	$c_3 - c_2$
231	0	c_2	$c_3 - c_2$
312	0	0	c_3
321	0	0	c_3
	$\frac{c_1}{3}$	$\frac{c_1}{3} + \frac{c_2 - c_1}{2}$	$c_3 - \frac{c_2}{2} - \frac{c_1}{6}$

Remark

The first player uses only one km. He equally shares the cost c_1 with the other players. The second km has a marginal cost of $c_2 - c_1$, equally shared by the players using it, the rest is paid by the player, the only one using the third km

Interesting properties for a solution

Cooperative
games (2)

Roberto
Lucchetti

The nucleolus

The Shapley
value

Power indices

Semivalues

The Shapley value, among others, satisfies the following important properties:

- ❶ For every $v \in \mathcal{G}(N)$

$$\sum_{i \in N} \sigma_i(v) = v(N)$$

- ❷ Let $v \in \mathcal{G}(N)$ be a game with the following property, for players i, j : for every A not containing i, j , $v(A \cup \{i\}) = v(A \cup \{j\})$.
Then

$$\sigma_i(v) = \sigma_j(v)$$

- ❸ Let $v \in \mathcal{G}(N)$ and $i \in N$ be such that $v(A) = v(A \cup \{i\})$ for all A . Then

$$\sigma_i(v) = 0$$

- ❹ for every $v, w \in \mathcal{G}(N)$, $\sigma(v + w) = \sigma(v) + \sigma(w)$

Comments

Cooperative
games (2)

Roberto
Lucchetti

The nucleolus

The Shapley
value

Power indices

Semivalues

- ▶ Property 1 is **efficiency**
- ▶ Property 2 is **symmetry**: symmetric players must take the same
- ▶ Property 3 is **Null player property**: a player contributing nothing to any coalition must have nothing
- ▶ Property 4 is **additivity**

The Shapley value for simple games

Cooperative
games (2)

Roberto
Lucchetti

The nucleolus

The Shapley
value

Power indices

Semivalues

In the case of the simple games, the Shapley value becomes

$$\sigma_i(v) = \sum_{A \in \mathcal{A}_i} \frac{(a-1)!(n-a)!}{n!},$$

where \mathcal{A}_i is the set of the coalitions A such that

- $i \in A$
- A is winning
- $A \setminus \{i\}$ is not winning

Weighted voting doesn't work

Cooperative
games (2)

Roberto
Lucchetti

The nucleolus

The Shapley
value

Power indices

Semivalues

In a stock company A owns the 10% of the stock, B the 20%, C the 30% and finally D the 40%.

The Shapley value of the associated simple game is

$$\sigma = \left(\frac{1}{12}, \frac{3}{12}, \frac{3}{12}, \frac{5}{12} \right).$$

Note the difference with the proportional ownership:

$$\left(\frac{1}{10}, \frac{2}{10}, \frac{3}{10}, \frac{4}{10} \right).$$

Now suppose there is a little exchange between B and D providing the new situation: A owns the 10% of the stock, B the 21%, C the 30% and finally D the 39%.

The new Shapley becomes

$$\left(0, \frac{1}{3}, \frac{1}{3}, \frac{1}{3} \right).$$

Probabilistic power indices

Cooperative
games (2)

Roberto
Lucchetti

The nucleolus

The Shapley
value

Power indices

Semivalues

In simple games the Shapley value assumes also the meaning of measuring **the fraction of power of every player**. To measure the relative power of the players in a simple game, the efficiency requirement is not anymore mandatory, and the way coalitions could form can be different from the case of the Shapley value

Definition

A **probabilistic power index** ψ on the set of simple games is

$$\psi_i(v) = \sum_{S \in 2^{N \setminus \{i\}}} p_i(S) m_i(v, S)$$

where p_i is a probability measure on $2^{N \setminus \{i\}}$

Remark

Remember: $m_i(v, S) = v(S \cup \{i\}) - v(S)$

Semivalues

Cooperative
games (2)

Roberto
Lucchetti

The nucleolus

The Shapley
value

Power indices

Semivalues

Definition

A probabilistic power index ψ on the set of simple games is a **semivalue** if there exists a vector (p_0, \dots, p_{n-1}) such that

$$\psi_i(v) = \sum_{S \in 2^{N \setminus \{i\}}} p_s m_i(v, S)$$

$p_i(S) = p_s$, i.e. the coefficient does not depend from player i and depends from the coalition S only through its cardinality s .

Remark

Since the index is probabilistic, the two conditions must hold

- $p_s \geq 0$
- $\sum_{n=0}^{n-1} \binom{n-1}{s} p_s = 1$

If $p_s > 0$ for all s , the semivalue is called **regular**

Examples

Cooperative
games (2)

Roberto
Lucchetti

The nucleolus

The Shapley
value

Power indices

Semivalues

These are examples of semivalues

- the Shapley value
- the **Banzhaf** value

$$\beta_i(v) = \sum_{S \in 2^{N \setminus \{i\}}} \frac{1}{2^{n-1}} (v(S \cup \{i\}) - v(S)).$$

- the **Binomial values**: $p_s = q^s(1 - q)^{n-s-1}$, for every $0 < q < 1$
- the **marginal value**, $p_s = 0$ for $s = 0, \dots, n-2$: $p_{n-1} = 1$
- the **dictatorial value** $p_s = 0$ for $s = 1, \dots, n-1$: $p_0 = 1$

The U.N. security council

Cooperative games (2)

Roberto
Lucchetti

The nucleolus

The Shapley
value

Power indices

Semivalues

Example

Let $N = \{1, \dots, 15\}$. The permanent members 1, . . . 5 are veto players. A resolution passes provided it gets at least 9 votes, including the five votes of the permanent members

- ▶ Let i be a player which is no veto. His marginal value is 1 if and only if it enters a coalition A such that $a = 8$ and A contains the 5 veto players. Then

$$\sigma_i = \frac{8! \cdot 6! \cdot 9 \cdot 8 \cdot 7}{15! \cdot 3 \cdot 2} \simeq 0.0018648$$

- ▶ The power of the veto player j can be calculated by difference and symmetry. The result is $\sigma_j \simeq 0,1962704$

Calculating Banzhaf power index

- ▶ Let i be a player which is no veto. Then

$$\beta_i = \frac{1}{2^{14}} \binom{9}{3} = \frac{21}{2^{12}} \simeq 0.005127$$

- ▶ Let j be a veto player. Then

$$\beta_j = \frac{1}{2^{14}} \left(\binom{10}{4} + \dots + \binom{10}{10} \right) = \frac{1}{2^{14}} \left(2^{10} - \sum_{k=0}^3 \binom{10}{k} \right) = \frac{53}{2^{10}} \simeq 0.0517578$$

Remark

- ▶ The ratio $\frac{\sigma_i}{\sigma_j} \simeq 105.25$

- ▶ The ratio $\frac{\beta_i}{\beta_j} \simeq 10.0951$